

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 July 2001 (19.07.2001)

PCT

(10) International Publication Number
WO 01/51042 A2

(51) International Patent Classification⁷: **A61K 31/00**

SIEGEL, Richard, Lewis [US/US]; Pfizer Inc., 235 East 42nd Street, New York, NY 10017 (US).

(21) International Application Number: **PCT/IB01/00018**

(22) International Filing Date: 11 January 2001 (11.01.2001)

(74) Agents: **WOOD, David, J. et al.**; Pfizer Limited, Ramsgate Road, Sandwich, Kent CT13 9NJ (GB).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0000561.1 11 January 2000 (11.01.2000) GB

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(71) Applicant (*for GB only*): **PFIZER LIMITED** [GB/GB]; Ramsgate Road, Sandwich, Kent CT13 9NJ (GB).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except GB, US*): **PFIZER INC.** [US/US]; 235 East 42nd Street, New York, NY 10017 (US).

(71) Applicant and
(72) Inventor: **WOOD, Ralph, E.** [US/US]; R.D. #2, Box 193A, Moundsville, WV 26041 (US).

Published:
— without international search report and to be republished upon receipt of that report

(72) Inventors; and
(75) Inventors/Applicants (*for US only*): **DAVIES, Michael, John** [GB/GB]; Pfizer Global Research and Development, Ramsgate Road, Sandwich, Kent CT13 9NJ (GB).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/51042 A2

(54) Title: **TREATMENT OF DIABETIC ULCERS**

(57) Abstract: This invention relates to the use of cyclic guanosine 3', 5'-monophosphate phosphodiesterase type five (cGMP PDE5) inhibitors, including in particular the compound sildenafil, for the treatment of diabetic ulcers, particularly diabetic foot ulcers.

TREATMENT OF DIABETIC ULCERS

This invention relates to the use of cyclic guanosine 3', 5'-monophosphate type five (cGMP PDE5) inhibitors, including in particular the compound sildenafil, for
5 the treatment of diabetic ulcers including in particular the treatment of chronic dermal ulcers such as diabetic foot ulcers.

Diabetes is a serious chronic disease. The 1996 global diabetes prevalence of 120 million is predicted to more than double to 250 million by the year 2025 due
10 to increasing age, obesity, sedentary lifestyle and changing dietary patterns. Though many serious and costly complications affect individuals with diabetes, such as heart disease, kidney failure and blindness, foot complications take the greatest toll: 40-70% of all lower extremity amputations are related to diabetes mellitus. Indeed, 85% of all diabetes-related lower extremity amputations are
15 preceded by a foot ulcer.

Patients with diabetes mellitus are known to be at increased risk of developing chronic dermal ulcers such as an ulcer on the foot in the presence of established long-term complications of the disease. Ulceration occurs as the result of
20 impaired nerve function (neuropathy) and/or in the presence of ischaemia.

Local tissue ischaemia is a major contributing factor to diabetic ulceration. As well as large vessel disease, patients with diabetes suffer additional threat to their skin perfusion by involvement of the non-conduit arteries in the process of
25 atherosclerosis and, perhaps more importantly, impairment of the microcirculatory control mechanisms, so-called small vessel disease. Under normal circumstances, blood flow increases in response to injury to facilitate healing. In the presence of small vessel disease (and ischaemia) this response is significantly blunted and this, together with the tendency to thrombosis in the
30 microcirculation during low flow, is probably important in ulcer pathogenesis.

Neuropathy is also a major complication of diabetes mellitus, with no well-established therapy for either its symptomatic treatment or for prevention of progressive decline in nerve function. Estimates of the prevalence of neuropathy in diabetes vary widely (5 to 80%), largely due to the wide variety of definitions and clinical descriptions of neuropathy. Nevertheless, prevalence rates in the order of 20% have been recorded in both hospital and community-based studies in the UK.

The effect of the neuropathy is complex. However, loss of sensory information from the foot undoubtedly contributes to abnormal and prolonged pressure on the areas of the foot (sensory neuropathy). Motor-neuropathy causes deformity, further increasing pressure loading. In autonomic neuropathy, loss of innervation of the sweat glands results in dry skin which cracks allowing foci for infection. Autonomic dysfunction contributes further by altering the distribution of micro-circulatory blood flow, directing the blood flow through shunts and away from the nutritive skin capillaries. It is the interplay of these factors and foot trauma that result in the skin breakdown.

The mechanism leading to nerve damage in diabetes is not yet resolved but is almost certainly multifactorial, with genetic predisposition, metabolic and vascular abnormalities, and lack of perturbation of growth factors, implicated. The response of the peripheral nervous system to the metabolic insults received in diabetes does not seem to differ between type 1 and type 2 diabetes, suggesting the likelihood of similar clinical response to therapies in the two primary forms of the disease. There appears to be a number of susceptibility factors, as yet unknown, for the development of neuropathy, which operate in the presence of hyperglycaemia.

According to the invention, there is provided a method of treating a patient with diabetic ulcers which comprises treating the patient with an effective amount of a cGMP PDE5 inhibitor, or a pharmaceutical composition thereof.

The invention also provides for the use of a cGMP PDE5 inhibitor for the manufacture of a composition for the treatment of diabetic ulcers.

5 The invention is of particular value for treating patients with diabetic foot ulceration.

A number of potent and selective cGMP PDE5 inhibitors are now known and their activity can be readily determined by in-vitro screening against cGMP PDE enzymes from a number of sources, in accordance with published procedures.

10 Thus for example a number of pyrazolopyrimidinone compounds are described as cGMP PDE5 inhibitors in our European patents nos. 0463756 and 0526004; while further compounds are described in International patent applications nos. WO 93/12095; WO 94/05661; WO 94/00453; WO 95/19978 and WO 98/49166.

15 Preferred cGMP-PDE5 inhibitors for use in the present invention include:

5-(2-ethoxy-5-morpholinoacetylphenyl)-1-methyl-3-n-propyl-1,6-dihydro-7H-pyrazolo[4, 3-d]pyrimidin-7-one;

20 5-(5-morpholinoacetyl-2-n-propoxyphenyl)-1-methyl-3-n-propyl-1,6-dihydro-7H-pyrazolo[4, 3-d]pyrimidin-7-one;

5-[2-ethoxy-5-(4-methyl-1-piperazin-1-yl-sulphonyl)-phenyl]-1,6-dihydro-1-methyl-3-propylpyrazolo[4, 3-d]pyrimidin-7-one;

5-[2-allyloxy-5-(4-methyl-1-piperazinylsulphonyl)-phenyl]-1-methyl-3-n-propyl-1,6-dihydro-7H-pyrazolo[4, 3-d]pyrimidin-7-one;

25 5-(2-ethoxy-5-[4-(2-propyl)-1-piperazinyl-sulphonyl]phenyl)-1-methyl-3-n-propyl-1,6-dihydro-7H-pyrazolo[4, 3-d]pyrimidin-7-one;

5-(2-ethoxy-5[4-(2-hydroxyethyl-1-piperazinylsulphonyl]phenyl)-1-methyl-3-n-propyl-1, 6-dihydro-7H-pyrazolo [4, 3-d]pyrimidin-7-one;

30 5-(5-[4-(2-hydroxyethyl)-1-piperazinylsulphonyl]-2-n-propoxyphenyl)-1-methyl-3-n-propyl-1, 6-dihydro-7H-pyrazolo[4, 3-d]pyrimidin-7-one;

5-[2-ethoxy-5-(4-methyl-1-piperazinylcarbonyl)-phenyl]-1-methyl-3-n-propyl-1, 6-dihydro-7H-pyrazolo[4, 3-d]pyrimidin-7-one;

5-[2-ethoxy-5-(1-methyl-2-imidazolyl)phenyl]-1-methyl-3-n-propyl-1, 6-dihydro-7H-pyrazolo[4, 3-d]pyrimidin-7-one,
3-ethyl-5-[5-[4-ethylpiperazin-1-yl)sulphonyl]-2-(2-methoxyethoxy)pyrid-3-yl]-2-(2-pyridylmethyl)-6,7-dihydro-2H-pyrazolo[4,3-d]pyrimidin-7-one,
5 3-ethyl-5-[5-(4-ethylpiperazin-1-ylsulphonyl)-2-n-propoxyphenyl]-2-(pyridin-2-yl)methyl-2,6-dihydro-7H-pyrimidin-7-one, and
5-[2-ethoxy-5-(4-ethylpiperazin-1-ylsulphonyl)pyridin-3-yl]-3-ethyl-2-[2-methoxyethyl]-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one, also known
as 1-[6-ethoxy-5-[3-ethyl-6,7-dihydro-2-(2-methoxyethyl)-7-oxo-2H-
10 pyrazolo[4,3-d]pyrimidin-5-yl]-3-pyridylsulphonyl]-4-ethylpiperazine (see
Examples hereinafter for preparation).

Most particular preferred is the compound 5-[2-ethoxy-5-(4-methylpiperazin-1-ylsulphonyl)-phenyl]-1, 6-dihydro-1-methyl-3-propylpyrazolo [4,3-d] pyrimidine-7-
15 one (sildenafil), and pharmaceutically acceptable salts thereof; including the citrate salt.

The suitability of any particular cGMP PDE5 inhibitor can be readily determined by evaluation of its potency and selectivity using literature methods followed by
20 evaluation of its toxicity, absorption, metabolism, pharmacokinetics, etc in accordance with standard pharmaceutical practice.

Preferably, the cGMP PDE5 inhibitors have an IC₅₀ for PDE5 at less than 100 nanomolar, more preferably, at less than 50 nanomolar, more preferably still at
25 less than 10 nanomolar.

IC₅₀ values for the cGMP PDE5 inhibitors may be determined using established literature methodology, for example as described in EP0463756-B1 and EP0526004-A1.

30

Preferably the cGMP PDE5 inhibitors used in the invention are selective for the PDE5 enzyme. Preferably they are selective over PDE3, more preferably over

PDE3 and PDE4. Preferably, the cGMP PDE5 inhibitors of the invention have a selectivity ratio greater than 100 more preferably greater than 300, over PDE3 and more preferably over PDE3 and PDE4.

- 5 Selectivity ratios may readily be determined by the skilled person. IC50 values for the PDE3 and PDE4 enzyme may be determined using established literature methodology, see S A Ballard *et al*, Journal of Urology, 1998, vol. 159, pages 2164-2171.
- 10 The cGMP PDE5 inhibitors can be administered alone but, in human therapy will generally be administered in admixture with a suitable pharmaceutical excipient diluent or carrier selected with regard to the intended route of administration and standard pharmaceutical practice.
- 15 For example, the cGMP PDE5 inhibitors can be administered orally, buccally or sublingually in the form of tablets, capsules, ovules, elixirs, solutions or suspensions, which may contain flavouring or colouring agents, for immediate-, delayed-, modified-, or controlled-release applications.
- 20 Surprisingly, PDE5 inhibitors treat diabetic ulcers systemically. Accordingly, oral administration is a preferred route.

- Such tablets may contain excipients such as microcrystalline cellulose, lactose, sodium citrate, calcium carbonate, dibasic calcium phosphate and glycine,
- 25 disintegrants such as starch (preferably corn, potato or tapioca starch), sodium starch glycollate, croscarmellose sodium and certain complex silicates, and granulation binders such as polyvinylpyrrolidone, hydroxypropylmethyl cellulose, hydroxypropylcellulose, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, stearic acid, glyceryl behenate and talc
- 30 may be included.

Solid compositions of a similar type may also be employed as fillers in gelatin capsules. Preferred excipients in this regard include lactose, starch, a cellulose, milk sugar or high molecular weight polyethylene glycols. For aqueous suspensions and/or elixirs, the cGMP PDE5 inhibitors of the invention may be
5 combined with various sweetening or flavouring agents, colouring matter or dyes, with emulsifying and/or suspending agents and with diluents such as water, ethanol, propylene glycol and glycerin, and combinations thereof.

The cGMP PDE5 inhibitors can also be administered parenterally, for example,
10 intravenously, intra-arterially, intraperitoneally, intramuscularly or subcutaneously, or they may be administered by infusion techniques. For such parenteral administration they are best used in the form of a sterile aqueous solution which may contain other substances, for example, enough salts or glucose to make the solution isotonic with blood. The aqueous solutions should
15 be suitably buffered (preferably to a pH of from 3 to 9), if necessary. The preparation of suitable parenteral formulations under sterile conditions is readily accomplished by standard pharmaceutical techniques well-known to those skilled in the art.

20 The dosage of cGMP PDE5 inhibitor in such formulations will depend on its potency, but can be expected to be in the range of from 1 to 500 mg for administration up to three times a day. For oral and parenteral administration to human patients, the daily dosage level of the cGMP PDE5 inhibitor will usually be from 5 to 500 mg (in single or divided doses). In the case of sildenafil, a preferred
25 dose is in the range 10 to 100 mg (e.g. 10, 25, 50 and 100 mg) which can be administered once, twice or three times a day (preferably once). However the precise dose will be as determined by the prescribing physician and will depend on the age and weight of the patient and severity of the symptoms.

30 Thus, for example, tablets or capsules of the cGMP PDE5 inhibitor may contain from 5 to 250 mg (e.g. 10 to 100 mg) of active compound for administration singly or two or more at a time, as appropriate. The physician in any event will

determine the actual dosage which will be most suitable for any individual patient and it will vary with the age, weight and response of the particular patient. The above dosages are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited and such
5 are within the scope of this invention.

The cGMP PDE5 inhibitors can also be administered intranasally or by inhalation and are conveniently delivered in the form of a dry powder inhaler or an aerosol spray presentation from a pressurised container, pump, spray or nebuliser with
10 the use of a suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, a hydrofluoroalkane such as 1,1,1,2-tetrafluoroethane or 1,1,1,2,3,3,3-heptafluoropropane, carbon dioxide or other suitable gas. In the case of a pressurised aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. The pressurised
15 container, pump, spray or nebuliser may contain a solution or suspension of the cGMP PDE5 inhibitor, e.g. using a mixture of ethanol and the propellant as the solvent, which may additionally contain a lubricant, e.g. sorbitan trioleate. Capsules and cartridges (made, for example, from gelatin) for use in an inhaler or insufflator may be formulated to contain a powder mix of the cGMP PDE5
20 inhibitor and a suitable powder base such as lactose or starch.

Aerosol or dry powder formulations are preferably arranged so that each metered dose or "puff" contains from 1 to 50 mg of the cGMP PDE5 inhibitor, for delivery to the patient. The overall daily dose with an aerosol will be in the range of from
25 1 to 50 mg which may be administered in a single dose or, more usually, in divided doses throughout the day.

Alternatively, the cGMP PDE5 inhibitors can be administered in the form of a suppository or pessary.

30

The cGMP PDE5 inhibitor may be applied topically in the form of a gel, hydrogel, lotion, solution, cream, ointment or dusting powder. The cGMP PDE5 inhibitors

may also be dermally or transdermally administered; for example, by the use of a skin patch.

Since diabetic ulcers occur on the skin surface, topical administration is a
5 preferred route of administration.

For application topically to the skin, the cGMP PDE5 inhibitors can be formulated as a suitable ointment containing the inhibitor suspended or dissolved in, for example, a mixture with one or more of the following: mineral oil, liquid
10 petrolatum, white petrolatum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, they can be formulated as a suitable lotion or cream, suspended or dissolved in, for example, a mixture of one or more of the following: mineral oil, sorbitan monostearate, a polyethylene glycol, liquid paraffin, polysorbate 60, cetyl esters
15 wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

The cGMP PDE5 inhibitors may also be used in combination with a cyclodextrin. Cyclodextrins are known to form inclusion and non-inclusion complexes with drug molecules. Formation of a drug-cyclodextrin complex may modify the solubility,
20 dissolution rate, bioavailability and/or stability property of a drug molecule. Drug-cyclodextrin complexes are generally useful for most dosage forms and administration routes. As an alternative to direct complexation with the drug the cyclodextrin may be used as an auxiliary additive, e.g. as a carrier, diluent or solubiliser. Alpha-, beta- and gamma-cyclodextrins are most commonly used and
25 suitable examples are described in WO-A-91/11172, WO-A-94/02518 and WO-A-98/55148.

Generally, in humans, oral administration of the cGMP PDE5 inhibitors is the preferred route, being the most convenient. In circumstances where the recipient
30 suffers from a swallowing disorder or from impairment of drug absorption after oral administration, the drug may be administered parenterally, sublingually or buccally.

It is to be appreciated that all references herein to treatment include curative, palliative and prophylactic treatment.

- 5 The following formulation examples are illustrative only and are not intended to limit the scope of the invention . Active ingredient means a cGMP PDE5 inhibitor.

Formulation 1:

A tablet is prepared using the following ingredients :

- 10 Sildenafil citrate (50 mg) is blended with cellulose (microcrystalline), silicon dioxide, stearic acid (fumed) and the mixture is compressed to form tablets.

Formulation 2 :

- 15 An intravenous formulation may be prepared by combining active ingredient (100 mg) with isotonic saline (1000 ml).


The efficacy of the cGMP PDE5 inhibitors in treating neuropathy in human patients is illustrated by the following anecdotal report.

- 20 The patient was an insulin-dependent diabetic who had been suffering from erectile dysfunction and who subsequently developed a diabetic foot ulcer. During treatment of his erectile dysfunction with sildenafil it was noted that his foot ulcer, which he had had for approximately two years, was healing. The patient had been through various courses of treatment with minimal results. He
25 had seen vascular surgeons, podiatrists and had visited wound care clinics. Ultimately, he was hospitalised for approximately one month on IV antibiotics etc, and the threat was very real that the patient was going to require a below-the-knee amputation. At times the ulcer would appear to be healing only to re-occur, returning to its pre-treatment size and depth. Once Sildenafil treatment had
30 begun for his erectile dysfunction, it was noted that the ulcer was decreasing in size. The patient was instructed to continue taking 50 mg of Sildenafil once a day. This treatment resulted in complete resolution of the diabetic foot ulcer

within one month. The patient has continued on this same treatment for the past two years without re-occurrence.

Examples

- 5 2-(Methoxyethyl)-5-[2-ethoxy-5-(4-ethylpiperazin-1-ylsulphonyl)pyridin-3-yl]-3-ethyl-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one




- A mixture of the product from stage i) below (0.75mmol), potassium
bis(trimethylsilyl)amide (298mg, 1.50mmol) and ethyl acetate (73 microlitres,
10 0.75mmol) in ethanol (10ml) was heated at 120°C in a sealed vessel for 12
hours. The cooled mixture was partitioned between ethyl acetate and aqueous
sodium bicarbonate solution, and the layers separated. The organic phase was
dried (MgSO₄), and evaporated under reduced pressure. The crude product was
purified by column chromatography on silica gel using dichloromethane:methanol
15 (98:2) as eluant to afford the title compound, 164mg; Found : C, 53.18; H, 6.48;
N, 18.14; C₂₃H₃₃N₇O₅S; 0.20C₂H₅CO₂CH₃ requires C, 53.21; H, 6.49; N, 18.25%; δ
(CDCl₃) : 1.04 (3H, t), 1.40 (3H, t), 1.58 (3H, t), 2.41 (2H, q), 2.57 (4H, m), 3.08
(2H, q), 3.14 (4H, m), 3.30 (3H, s), 3.92 (2H, t), 4.46 (2H, t), 4.75 (2H, q), 8.62
(1H, d), 9.04 (1H, d), 10.61 (1H, s); LRMS : m/z 520 (M+1)⁺; mp 161-162°C.

20

Preparation of Starting Materials


- a) Pyridine-2-amino-5-sulphonic acid

11




2-Aminopyridine (80g, 0.85mol) was added portionwise over 30 minutes to oleum (320g) and the resulting solution heated at 140°C for 4 hours. On cooling, the reaction was poured onto ice (200g) and the mixture stirred in an ice/salt bath for a further 2 hours. The resulting suspension was filtered, the solid washed with ice water (200ml) and cold IMS (200ml) and dried under suction to afford the title compound as a solid, 111.3g; LRMS : m/z 175 (M+1)⁺.

10 b) Pyridine-2-amino-3-bromo-5-sulphonic acid



Bromine (99g, 0.62mol) was added dropwise over an hour, to a hot solution of the product from stage a) (108g, 0.62mol) in water (600ml) so as to maintain a steady reflux. Once the addition was complete the reaction was cooled and the resulting mixture filtered. The solid was washed with water and dried under suction to afford the title compound, 53.4g; δ (DMSO-d₆, 300MHz): 8.08 (1H, s), 8.14 (1H, s); LRMS : m/z 253 (M)⁺.


20 c) Pyridine-3-bromo-2-chloro-5-sulphonyl chloride



A solution of sodium nitrite (7.6g, 110.0mmol) in water (30ml) was added dropwise to an ice-cooled solution of the product from stage b) (25.3g,

100.0mmol) in aqueous hydrochloric acid (115ml, 20%), so as to maintain the temperature below 6°C. The reaction was stirred for 30 minutes at 0°C and for a further hour at room temperature. The reaction mixture was evaporated under reduced pressure and the residue dried under vacuum at 70°C for 72 hours. A mixture of this solid, phosphorus pentachloride (30.0g, 144mmol) and phosphorus oxychloride (1ml, 10.8mmol) was heated at 125°C for 3 hours, and then cooled. The reaction mixture was poured onto ice (100g) and the resulting solid filtered, and washed with water. The product was dissolved in dichloromethane, dried (MgSO₄), and evaporated under reduced pressure to afford the title compound as a yellow solid, 26.58g; δ (CDCl₃, 300MHz) : 8.46 (1H, s), 8.92 (1H, s).


d) 3-Bromo-2-chloro-5-(4-ethylpiperazin-1-ylsulphonyl)pyridine



A solution of 1-ethylpiperazine (11.3ml, 89.0mmol) and triethylamine (12.5ml, 89.0mmol) in dichloromethane (150ml) was added dropwise to an ice-cooled solution of the product from stage c) (23.0g, 79.0mmol) in dichloromethane (150ml) and the reaction stirred at 0°C for an hour. The reaction mixture was concentrated under reduced pressure and the residual brown oil was purified by column chromatography on silica gel, using an elution gradient of dichloromethane:methanol (99:1 to 97:3) to afford the title compound as an orange solid, 14.5g; δ (CDCl₃, 300MHz) : 1.05 (3H, t), 2.42 (2H, q), 2.55 (4H, m), 3.12 (4H, m), 8.24 (1H, s), 8.67 (1H, s).


e) 3-Bromo-2-ethoxy-5-(4-ethylpiperazin-1-ylsulphonyl)pyridine

13



A mixture of the product from stage d) (6.60g, 17.9mmol) and sodium ethoxide (6.09g, 89.55mmol) in ethanol (100ml) was heated under reflux for 18 hours, then cooled. The reaction mixture was concentrated under reduced pressure, the residue partitioned between water (100ml) and ethyl acetate (100ml), and the layers separated. The aqueous phase was extracted with ethyl acetate (2x100ml), the combined organic solutions dried (MgSO₄) and evaporated under reduced pressure to afford the title compound as a brown solid, 6.41g; Found : C, 41.27; H, 5.33; N, 11.11. C₁₃H₂₀BrN₃O₃S requires C, 41.35; H, 5.28; N, 10.99%; δ (CDCl₃, 300MHz) : 1.06 (3H, t), 1.48 (3H, t), 2.42 (2H, q), 2.56 (4H, m), 3.09 (4H, m), 4.54 (2H, q), 8.10 (1H, s), 8.46 (1H, s); LRMS : m/z 378, 380 (M+1)⁺.


f) Pyridine 2-ethoxy-5-(4-ethylpiperazin-1-ylsulphonyl)-3-carboxylic acid ethyl ester



A mixture of the product from stage e) (6.40g, 16.92mmol), triethylamine (12ml, 86.1mmol), and palladium (0) tris(triphenylphosphine) in ethanol (60ml) was heated at 100°C and 200 psi, under a carbon monoxide atmosphere, for 18 hours, then cooled. The reaction mixture was evaporated under reduced pressure and the residue purified by column

chromatography on silica gel, using an elution gradient of dichloromethane:methanol (100:0 to 97:3) to afford the title compound as an orange oil, 6.2g; δ (CDCl_3 , 300MHz) : 1.02 (3H, t), 1.39 (3H, t), 1.45 (3H, t), 2.40 (2H, q), 2.54 (4H, m), 3.08 (4H, m), 4.38 (2H, q), 4.55 (2H, q), 8.37 (1H, s), 8.62 (1H, s); LRMS : m/z 372 (M+1)⁺.


g) Pyridine 2-ethoxy-5-(4-ethylpiperazin-1-ylsulphonyl)-3-carboxylic acid



A mixture of the product from stage f) (4.96g, 13.35mmol) and aqueous sodium hydroxide solution (25ml, 2N, 50.0mmol) in ethanol (25ml) was stirred at room temperature for 2 hours. The reaction mixture was concentrated under reduced pressure to half its volume, washed with ether and acidified to pH 5 using 4N hydrochloric acid. The aqueous solution was extracted with dichloromethane (3x30ml), the combined organic extracts dried (MgSO_4) and evaporated under reduced pressure to afford the title compound as a tan coloured solid, 4.02g; δ (DMSO-d_6 , 300MHz) : 1.18 (3H, t), 1.37 (3H, t), 3.08 (2H, q), 3.17-3.35 (8H, m), 4.52 (2H, q), 8.30 (1H, s), 8.70 (1H, s).


h) 4-[2-Ethoxy-5-(4-ethylpiperazin-1-ylsulphonyl)pyridin-3-ylcarboxamido]-1H-3-ethylpyrazole-5-carboxamide

15



A solution of 4-amino-3-ethyl-1H-pyrazole-5-carboxamide (WO 9849166) (9.2g, 59.8mmol) in N,N-dimethylformamide (60ml) was added to a solution of the product from stage g) (21.7g, 62.9mmol), 1-hydroxybenzotriazole hydrate (10.1g, 66.0mmol) and triethylamine (13.15ml, 94.3mmol) in dichloromethane (240ml). 1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (13.26g, 69.2mmol) was added and the reaction stirred at room temperature for 6 hours. The dichloromethane was removed under reduced pressure, the remaining solution poured into ethyl acetate (400ml), and this mixture washed with aqueous sodium bicarbonate solution (400ml). The resulting crystalline precipitate was filtered, washed with ethyl acetate and dried under vacuum, to afford the title compound, as a white powder, 22g; δ (CDCl₃+1 drop DMSO-d₆) 0.96 (3H, t), 1.18 (3H, t), 1.50 (3H, t), 2.25-2.56 (6H, m), 2.84 (2H, q), 3.00 (4H, m), 4.70 (2H, q), 5.60 (1H, br s), 6.78 (1H, br s), 8.56 (1H, d), 8.76 (1H, d), 10.59 (1H, s), 12.10-12.30 (1H, s); LRMS: m/z 480 (M+1)⁺.

i) 2-Methoxyethyl-4-[2-ethoxy-5-(4-ethylpiperazin-1-ylsulphonyl)pyridin-3-ylcarboxamido]-3-ethylpyrazole-5-carboxamide



1-Bromo-2-methoxyethane (1.72mmol) was added to a solution of the product from stage h) (750mg, 1.56mmol) and caesium carbonate (1.12g, 3.44mmol) in N,N-dimethylformamide (15ml) and the reaction stirred at 60° C for 18 hours. The cooled mixture was partitioned between water and ethyl acetate, and the layers separated. The organic layer was dried (MgSO₄), concentrated under reduced pressure and azeotroped with toluene to give a solid. This product was recrystallised from ether, to afford the title compound as a white solid.

Claims

- 1 A method of treating a patient suffering from diabetic ulcers which
comprises treating said patient with an effective amount of a cGMP PDE5
5 inhibitor, or a pharmaceutical composition thereof.
- 2 The use of a cGMP PDE5 inhibitor for the manufacture of a
pharmaceutical composition for the treatment of diabetic ulcers.
- 10 3 A method or use as claimed in claim 1 or claim 2 wherein the diabetic
ulcers are foot ulcers.
- 4 A method or use as claimed in claim 1 or claim 2 wherein the cGMP PDE5
inhibitor is sildenafil.
- 15

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 July 2001 (19.07.2001)

PCT

(10) International Publication Number
WO 01/51042 A3

- (51) International Patent Classification⁷: **A61K 31/505** **SIEGEL, Richard, Lewis** [US/US]; Pfizer Inc., 235 East 42nd Street, New York, NY 10017 (US).
- (21) International Application Number: PCT/IB01/00018
- (22) International Filing Date: 11 January 2001 (11.01.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
0000561.1 11 January 2000 (11.01.2000) GB
- (71) Applicant (for GB only): **PFIZER LIMITED** [GB/GB]; Ramsgate Road, Sandwich, Kent CT13 9NJ (GB).
- (71) Applicant (for all designated States except GB, US): **PFIZER INC.** [US/US]; 235 East 42nd Street, New York, NY 10017 (US).
- (71) Applicant and
- (72) Inventor: **WOOD, Ralph, E.** [US/US]; R.D. #2, Box 193A, Moundsville, WV 26041 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **DAVIES, Michael, John** [GB/GB]; Pfizer Global Research and Development, Ramsgate Road, Sandwich, Kent CT13 9NJ (GB).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
- (88) Date of publication of the international search report:
10 January 2002
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/51042 A3

(54) Title: TREATMENT OF DIABETIC ULCERS

(57) Abstract: This invention relates to the use of cyclic guanosine 3', 5'-monophosphate phosphodiesterase type five (cGMP PDE5) inhibitors, including in particular the compound sildenafil, for the treatment of diabetic ulcers, particularly diabetic foot ulcers.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/18 01/00018

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/505

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, WPI Data, BIOSIS, MEDLINE, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 99 54333 A (BUNNAGE MARK EDWARD ;MATHIAS JOHN PAUL (GB); STREET STEPHEN DEREK) 28 October 1999 (1999-10-28) the whole document	1-4
A	BOULTON ANDREW J.M.: "End-stage complications of diabetic neuropathy: Foot ulceration" CANADIAN JOURNAL OF NEUROLOGICAL SCIENCES, vol. 21, no. 4, 1994, pages s18-s22, XP002164862 the whole document	1-4

☐ Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

6 April 2001

Date of mailing of the international search report

26/04/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Engl, B

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/18 01/00018

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9954333 A	28-10-1999	AU 2742599 A	08-11-1999
		BR 9909808 A	26-12-2000
		EP 1073658 A	07-02-2001
		NO 20005255 A	19-12-2000
<hr/>			